

**FY 2022 Summary of Wildlife Management Activities on
Travis County-managed Tracts of the Balcones Canyonlands Preserve**



White-tailed deer doe and fawn on the 35 acres tract of the Jollyville Unit.
Game camera photo.

**Travis County
Department of Transportation and Natural Resources
Natural Resources and Environmental Quality Division**



October 1, 2021 - September 30, 2022

Table of Contents

Introduction	3
Brown-headed cowbirds	4
Feral hogs.....	7
White-tailed deer	8
Literature Cited.....	9

List of Tables

Table 1. Summary of Brown-Headed Cowbird Trapping Effort, FY22, Travis County BCP, Texas.....	5
Table 2. Results of Brown-Headed Cowbird Trapping Effort, FY22, Travis County BCP, Texas.....	6
Table 3. Results of Feral Hog Trapping Effort, FY22, Travis County BCP, Texas.....	7
Table 4. Results of White-tailed Deer Population Survey, FY22, Travis County BCP, Texas.....	9

INTRODUCTION

The Balcones Canyonlands Preserve (BCP) was created in May 1996 to protect the endangered golden-cheeked warbler (*Setophaga chrysoparia*) and seven other federally endangered species. As specified in the Balcones Canyonlands Conservation Plan (USFWS 1996; hereafter BCCP), the City of Austin and Travis County developed and implemented a Land Management Plan (LMP) to provide guidelines for managing the BCP. The LMP Tier IIA-10 chapter (BCP 2007) describes the responsibilities and guidelines for wildlife management activities on the BCP, including controlling populations of non-native, exotic, feral, and nuisance animals.

Non-native, exotic, and feral animals are species that are not indigenous to an area and were typically brought in by human activity, either intentionally or unintentionally. Examples of non-native animals in Travis County include red-imported fire ants (*Solenopsis invicta*), tawny crazy ants (*Nylanderia fulva*), and feral hogs (*Sus scrofa*). These species may directly or indirectly negatively impact protected species in the form of habitat degradation, displacement, depredation, and reducing available foods. To meet the species protection and enhancement goals required under the terms and conditions of the BCCP, BCP land managers strive to remove non-native, exotic, and feral species, where feasible.

Nuisance animals on the BCP are native species that may present threats to public health and safety, property, habitat quality, or other natural resources due primarily to high population densities. Examples of nuisance animals on the BCP include brown-headed cowbirds (*Molothrus ater*) and white-tailed deer (*Odocoileus virginianus*). Management to control nuisance animals has been directed by the USFWS as part of the recovery plans for protected species (BCP 2007).

In Fiscal Year 2022 (01 October 2021 - 30 September 2022; hereafter FY22), wildlife management activities on Travis County-managed tracts of the BCP focused on five species of non-native, exotic, feral, and/or nuisance animals. Monitoring and control of brown-headed cowbirds, feral hogs, and white-tailed deer are described in their respective sections in this report. Monitoring and control of red-imported fire ants and tawny crazy ants are described in *Appendix I* and *Appendix W2* of the FY22 BCCP Annual Report.

BROWN-HEADED COWBIRDS

Introduction

Brown-headed cowbirds (hereafter cowbird) are known to parasitize the nests of the golden-cheeked warbler and the black-capped vireo (*Vireo atricapilla*), in addition to many other avian hosts (TPWD No Datea). Historically, there had been little documentation of nest parasitism by cowbirds on the BCP, except for auxiliary observations from BCP staff. The BCP (2007) LMP suggests that assessing and reducing cowbird parasitism should be a primary concern for the BCP and population management options should include physical removal of cowbirds or their eggs.

Cowbird trapping has been the subject of considerable research and management effort and is believed to be an important technique for the conservation of both species. At the U.S. Army Fort Hood military base, cowbird trapping has been credited for drastically reducing parasitism rates of black-capped vireos from 91% before cowbird management, to below 20% after a cowbird management program was implemented (Kostecke et al. 2005). Population modelling data from Fort Hood suggests that sustained cowbird management is required to maintain viable black-capped vireo populations (Wilsey et al. 2014). Cowbird management success has contributed to the recovery effort and subsequent 2018 delisting of the black-capped vireo, and continued management is necessary to maintain black-capped vireo populations (USFWS 2018).

Methods

Cowbird trapping in FY22 was conducted exclusively in the western half of the county. BCP and Travis County Parks staff operated four metal hybrid cowbird traps (6'x8') and one mega trap (16'x16'). To maximize trap effectiveness, traps on the BCP were distributed near golden-cheeked warbler and black-capped vireo breeding habitat. Hybrid traps were operated on the Nootsie tract (Nootsie), Reimers Ranch Park (Reimers Ranch 1), Kotrla tract (Kotrla-Upper) and on private property adjacent to the Kotrla tract (Kotrla-Lower, known as the Kotrla trap prior to FY22). The mega trap was operated at Hamilton Pool (Hamilton Pool Mega).

The Kotrla-Lower trap was opened on March 30, 2022 and baited with a group of live decoy birds (5 males, 2 females) acquired from a trapping effort on private land in Hays County. Additional cowbirds captured at the Kotrla-Lower trap were used to stock and open the other four cowbird traps during the trapping period. The addition

of the Kotrla-Upper trap on the Kotrla tract in FY22 was due to increased brood parasitism and presence of cowbirds in the nesting habitat of black-capped vireos in FY21.

Table 1 summarizes the FY22 cowbird trapping schedule and initial stocking numbers of decoy birds.

Table 1. Summary of Brown-Headed Cowbird Trapping Effort, FY22, Travis County BCP, Texas.

Trap	Date Opened	Date Closed	Initial Stocking Numbers (Male/Female)
Nootsie	April 1	June 16	4/2
Kotrla-Lower	March 30	June 17	5/2
Kotrla-Upper	April 6	June 17	6/1
Hamilton Pool Mega	April 6	June 17	8/1
Reimers Ranch 1	April 6	June 17	4/1

Traps were inspected and maintained daily throughout the trapping period and any non-target species were removed and released. If staff was unavailable to conduct trap checks, the trap was shut by placing a board over the opening. Water and feed (white proso millet and whole milo) were refreshed three times per week, and as needed. Plastic water baths and shade cloth were added to the traps to offset the impact of rising air temperatures on bird health and survivability. Some traps were reinforced with poultry fencing along the outer base edges to prevent attempted predation. The traps were shut down in June and cowbirds were euthanized by placing them in a container with carbon dioxide gas following Texas Parks and Wildlife Department (TPWD) protocol (TPWD No Datea).

Results

In FY22, a total of 162 individual cowbirds were captured (108 males, 50 females, 4 immature) across 371 days in operation. In FY21, a total of 220 individual cowbirds were captured (137 males, 83 females) across 300 days in operation. This is a decrease in total capture despite an increase of days in operation and the addition of a trap to a potential higher cowbird use area. Keeping the Reimers Ranch 2 trap out of operation in FY22 may have also added to the lower capture numbers.

While care is taken to seal outer base edges against predators, there were some instances of predation by Texas rat snakes (*Elaphe obsoleta lindheimeri*) gaining entry from the top opening at two of the traps. Additionally, three non-targets were

captured, including one cardinal (*Cardinalis cardinalis*) and two painted buntings (*Passerina ciris*) across three traps. All non-target species were released some distance away from the trap.

Table 2 summarizes cowbird capture at each trap by sex, days in operation, and trapping efficiency (or capture rate). Trap success was calculated by dividing the number of females captured by the number of days in operation (x 100).

Travis County Natural Resources aims to maintain a minimum goal of 20% trap efficiency for the program. The overall trap efficiency was 13.48% in FY22, a decrease from 27.33% in FY21, with only Reimers Ranch 1 exceeding the 20% threshold.

Table 2. Results of Brown-Headed Cowbird Trapping Effort, FY22, Travis County BCP, Texas.

Trap	Days in Operation	Males Captured	Females Captured	Immature Captured	Total Captured	Overall Trap Efficiency %
Nootsie	76	24	5	1	30	6.58
Kotrla-Lower	79	27	9	0	36	11.39
Kotrla-Upper	72	9	5	0	14	6.94
Hamilton Pool Mega	72	16	6	0	22	8.33
Reimers Ranch 1	72	32	25	3	60	34.72
Grand Totals	371	108	50	4	162	13.48

Only one of the seven black-capped vireo nests found on the Kotrla tract and one of the five nests found on Reimers Ranch Park/Pogue Springs Preserve showed signs of cowbird parasitism (cowbird eggs or nestling in nest). Both nests ultimately failed due to predation and possible weather events. Additionally, BCP staff did not observe any instances of golden-cheeked warbler adults feeding a cowbird fledgling while conducting enumeration surveys in FY22.

While only one trap (Reimers Ranch 1) was operated on Reimers Ranch Park due to staffing constraints in FY22, it had a capture rate three times higher than Kotrla-Lower, which has historically been one of the most productive traps due to its proximity to rural areas and livestock. Staff will continue to assess the needs and resources of cowbird management and future trapping efforts prior to the FY23 trapping period.

FERAL HOGS

Introduction

The BCP (2007) LMP directs land managers to control populations of feral hogs to minimize negative impacts to the native wildlife protected within the BCP. Feral hogs degrade wildlife habitats and may compete directly with native wildlife for resources. They are omnivorous, primarily consuming vegetation, mast, roots and tubers, and to a lesser degree a wide range of animal species, including invertebrates, reptiles, amphibians, small mammals, and birds (Bradley and Schmidly 2016; Hellgren 1993). Their rooting habits create severely disturbed areas, which may lead to a localized shift in plant succession and increase the potential for soil erosion (Bradley and Schmidly 2016). Feral hogs also destabilize wetland areas, springs, creeks and other riparian areas through excessive rooting and wallowing. Their threat to humans and livestock through the spread of disease has also been documented (Miller 1993; U.S. Department of Agriculture 1999). Potentially producing two litters a year, with an average litter size of four to eight piglets, feral hog numbers can expand rapidly if left unmanaged (Wildlife Services 2012).

Methods

Standard feral hog trapping operation for BCP staff includes setting up corral style stock panel traps baited with game feeders filled with dry feed corn. Staff can operate up to six traps at a time. Each trap utilizes a wireless camera system that sends real-time images to a trapper's cell phone. The trapper monitors feral hog activity inside the trap and chooses when to remotely close the trap door. The trapper strives to trap the entire sounder at the same time. Trapped feral hogs are humanely dispatched, and carcasses are composted on the BCP.

Results

In FY22, a total of 17 feral hogs were trapped and removed from the BCP, while management efforts in FY21 resulted in the removal of 39 feral hogs. Table 3 summarizes the number of feral hogs harvested on the Travis County BCP in FY22 based on tract and sex.

Table 3. Results of Feral Hog Trapping Effort, FY22, Travis County BCP, Texas.

Tract	Sex (Male/Female)	Total
Ribelin	3/7	10
Sam Hamilton East	1/1	2
Hamilton Pool	1/4	5

Reduced trapping numbers in FY21 and FY22 suggest a decrease in the feral hog population after high levels of trapping in FY17, FY18, and FY19 resulting in 174, 127, and 114 feral hogs removed, respectively. Since implementing a feral hog monitoring and trapping system in FY17, BCP staff have observed a steady decline in feral hog field observations. As populations have declined, BCP staff have observed an increased hesitancy among feral hogs to enter traps, making it more difficult to round up the small remaining number of animals.

While full eradication of feral hogs from the BCP is impossible, staff will continue to monitor for damage and target any remaining feral hogs to stop further expansion onto the preserve.

WHITE-TAILED DEER

Introduction

The BCP (2007) LMP directs that white-tailed deer populations be monitored and maintained at a level that allows for successful recruitment of plant species that make up habitat supporting BCP protected species. Central Texas historically has some of the highest population densities of white-tailed deer in the United States (Richards 2001). Past research in Central Texas (Russell and Fowler 1999, Russell and Fowler 2002, Russell et al. 2001) indicates that little or no regeneration of vital habitat components may occur in areas with an over-population of deer. This generated an effort to design and implement a white-tailed deer population monitoring and control program for Travis County BCP properties as specified in the LMP Tier IIA-10.

To stay within carrying capacity, TPWD recommends population levels in the Texas Hill Country of one deer per 8-35 acres (TPWD No Date^b). Since a primary goal for the BCP is to maintain successful hardwood regeneration to provide long-term golden-cheeked warbler habitat, Travis County recommends a conservative density of one deer per 20-35 acres for the BCP.

Methods

BCP staff conducted nighttime spotlight deer surveys on the Kotrla Unit of the BCP and the Hamilton Pool Preserve/Reimers Ranch Park/Pogue Springs Preserve (hereafter HP/RR/Pogue Preserve). Additionally, camera surveys were operated as described in Oetgen et al. (2008) on the Jollyville Unit. This camera survey method

provides a more accurate population estimation of white-tailed deer in areas where there is a lack of infrastructure needed to perform nighttime spotlight surveys (Roberts et al. 2006).

Results

Deer population survey data gathered from August to October 2021 was used to estimate FY22 deer densities. Table 4 summarizes deer population data on the BCP collected from spotlight and camera surveys. Deer population densities on the BCP remained within the TPWD recommended population level but were slightly above Travis County's management objective of one deer per 20-35 acres.

Deer harvesting did not occur on the BCP in FY22. Survey results indicate that it will likely be beneficial to harvest deer in FY23, but a final decision will be made after the FY23 surveys are completed and reviewed.

Table 4. Results of White-tailed Deer Population Survey, FY22, Travis County BCP, Texas.

Location	Estimated Acres/Deer	Estimated Composition (Buck/Doe/Fawn)
Jollyville Unit*	15	12/57/2
Kotrla Unit**	19	20/21/8
HP/RR/Pogue Preserve	13	41/110/47

*Includes Snowden and Vireo Ridge tracts

**Includes Kotrla and Scott & White tracts

LITERATURE CITED

- Balcones Canyonlands Preserve. 2007. Balcones Canyonlands Preserve- Land Management Plan, Tier IIA-10, Animal Management. Unpublished document. August 2007. Travis County, Texas, USA.
<https://www.traviscountytx.gov/tnr/nr/bccp/reports>
- Bradley, R. D., and D. J. Schmidly. 2016. The mammals of Texas. University of Texas Press, Austin, USA.
- Hellgren, E. C. 1993. Biology of feral hogs (*Sus scrofa*) in Texas. Pages 50–58 in C. W. Hanselka and J. F. Cadenhead, editors. Feral swine: a compendium for resource managers: proceedings of a conference. Texas Agricultural Extension Service, San Angelo, USA.
- Kostecke, R. M., S. G. Summers, G. H. Eckrich, and D. A. Cimprich. 2005. Effects of brown-headed cowbird (*Molothrus ater*) removal on black-capped vireo (*Vireo atricapilla*) nest success and population growth at Fort Hood, Texas. Ornithological Monographs 57:28–37.
- Miller, J. E., 1993. A national perspective on feral swine. Pages 9–16 in C. W. Hanselka and J. F. Cadenhead, editors. Feral swine: a compendium for resource managers: proceedings of a conference. Texas Agricultural Extension Service, San Angelo, USA.
- Oetgen, J. G., B. C. Lambert, Jr., and J. D. Whiteside. 2008. Surveying white-tailed deer populations using infrared-triggered cameras.
https://tpwd.texas.gov/publications/pwdpubs/media/pwd_br_w7000_1444.pdf
- Richards, B. 2001. The deer dilemma: to fence or not to fence? Conference sponsored by Texas Parks and Wildlife Department. Reicher Ranch. June 15, 2001.
- Roberts, C. W., B. L. Pierce, A. W. Braden, R. R. Lopez, N. J. Silvy, P. A. Frank, D. Random, Jr., and A. Jara. 2006. Comparison of camera and road survey estimates for white-tailed deer. Journal of Wildlife Management 70:263–267.
- Russell, F. L., and N. L. Fowler. 1999. Rarity of oak saplings in savannas and woodlands of the eastern Edwards Plateau, Texas. The Southwestern Naturalist 44(1):31–41.

- Russell, F. L., and N. L. Fowler. 2002. Failure of adult recruitment in *Quercus buckleyi* populations on the eastern Edwards Plateau, Texas. *American Midland Naturalist* 148:201–217.
- Russell, F. L., D. B. Zippen, and N. L. Fowler. 2001. Effects of white-tailed deer (*Odocoileus virginianus*) on plants, plant populations and communities: a review. *American Midland Naturalist* 146:1–26.
- Texas Parks and Wildlife Department [TPWD]. No Datea. Trapping brown-headed cowbirds to control songbird nest parasitism. www.tpwd.state.tx.us/publications/pwdpubs/media/pwd_bk_w7000_1148.pdf
- Texas Parks and Wildlife Department [TPWD]. No Dateb. Managing habitat for white-tailed deer in the Hill Country area of Texas. https://tpwd.texas.gov/publications/pwdpubs/media/pwd_rp_w7000_0193.pdf
- U.S. Department of Agriculture [USDA]. 1999. Wild pigs: hidden dangers for farmers and hunters. Agriculture Information Bulletin No. 620. USDA. Riverdale, MD.
- U.S. Fish and Wildlife Service [USFWS]. 1996. Final Environmental Impact Statement/Habitat Conservation Plan for Proposed Issuance of a Permit to Allow Incidental Take of the Golden-cheeked Warbler, Black-capped Vireo, and six karst invertebrates in Travis County, Texas.
- U.S. Fish and Wildlife Service [USFWS]. 2018. “Endangered and Threatened Wildlife and Plants; Removing the Black Capped Vireo From the Federal List of Endangered and Threatened Wildlife.” 83 *Federal Register* 73, 16228-16242 (16 April 2018). <https://www.govinfo.gov/content/pkg/FR-2018-04-16/pdf/2018-07350.pdf>
- Wildlife Services. 2012. Managing feral hog damage. Texas A&M AgriLife Extension Service Wildlife Services. Pamphlet No. EL-1925. 03/2012. <https://cdn-ext.agnet.tamu.edu/wp-content/uploads/2019/12/EL1925-managing-feral-hog-damage.pdf>

Wilsey, C. B., J. J. Lawler, D. Cimprich, and N. H. Schumaker. 2014. Dependence of the endangered black-capped vireo on sustained cowbird management. *Conservation Biology* 28: 561–571.